

InVivoMAb anti-mouse TIGIT

Lot Specific Information

Lot Number:	Lot Specific*
Volume:	Lot Specific*
Concentration:	Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein:	Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	BE0274
Clone:	1G9
Isotype:	Mouse IgG1, κ
Recommended Isotype Control(s):	InVivoMAb mouse IgG1 isotype control, unknown specificity
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Immunogen:	Mouse TIGIT
Reported Applications:	<i>in vivo</i> TIGIT stimulation Flow cytometry
Formulation:	PBS, pH 7.0 Contains no stabilizers or preservatives
Endotoxin:	<2EU/mg (<0.002EU/μg) Determined by LAL gel clotting assay
Purity:	>95% Determined by SDS-PAGE
Sterility:	0.2 μm filtered
Production:	Purified from tissue culture supernatant in an animal free facility
Purification:	Protein G
RRID:	AB_2687797
Molecular Weight:	150 kDa

Description

The 1G9 monoclonal antibody reacts with mouse TIGIT (T cell immunoreceptor with Ig and ITIM domains). TIGIT is a 26 kDa, type I transmembrane protein and a member of the poliovirus receptor (PVR) family. TIGIT has been found to be expressed on follicular T helper cells in mice while in humans it's expressed by many T cell subsets including activated T cells, follicular T helper cells, memory T cells, and regulatory T cells as well as on NK cells. TIGIT can interact with certain members of the PVR and PVR-like families, including PVR, PVRL2, PVRL3, CD155, and CD112. TIGIT is thought to negatively regulate NK and T cell activation. Binding of TIGIT on T cells by dendritic cells results in their differentiation into a tolerogenic phenotype, with increased secretion of IL-10 and diminished production of IL-12. TIGIT knock-out mice are more susceptible to autoimmune disease.

Shelf-life and Storage

Store at the stock concentration at 4°C. **Do not freeze.**

All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at bxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit <https://bxcell.com/product/invivomab-anti-mouse-tigit/#references> or scan the QR code below.

Bio X Cell, Inc.

bxcell.com
1.866.787.3444
customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell



Bio X Cell, Inc.

bxcell.com
1.866.787.3444

customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell